- Example (Committees, LNp.5-6)

$(x-\mu)^2$	$x-\mu$	x	x^2	$f(x)$	$xf(x)$	$(x-\mu)^2 f(x)$	$x^2 f(x)$
4	-2	0	0	5/210	0/210	20/210	0/210
1	-1	1	1	50/210	50/210	50/210	50/210
0	0	2	4	100/210	200/210	0/210	400/210
1	1	3	9	50/210	150/210	50/210	450/210
4	2	4	16	5/210	20/210	20/210	80/210
		Totals		1	2 $E(X)$	2/3 $Var(X)$	14/3 $E(X^2)$

distribution; deviation of x from μ

So, $\mu = 2$, $\sigma^2 = 2/3$, and $\sigma = \sqrt{2/3}$



- Note.
  - μ_X and σ_X^2 only depends on f_X. They are *fixed* constants, not *random* numbers.
  - If X has units, then μ_X and σ_X have the same unit as X, and variance has unit squared.

X (random) cf. distribution (fixed)

acting like a scale of a ruler ⇒ standard deviation ⇒ high probability within $\mu \pm 2\sigma$, $\sigma^2=1, \pm 2$, $\sigma^2=100, \pm 20$

➢ Intuitive Interpretation of Variance → $E[(X-\mu_X)^2]$

- Variance is the weighted (權重 $f_X(x)$) average value of the squared deviation of X from μ_X.
- Variance is related to how the pmf is spread out



➢ Some properties of variance.

$E[(X-\mu_X)^2] = Var(X) \geq 0$

- The variance of a r.v. is always non-negative
- The only r.v. with variance equal to zero is a r.v. which can only take on a single value (μ_X).



It's actually not random

prove for discrete r.v.'s (but it's also true for other types of r.v.'s)

$\sum_{x \in \mathcal{X}} (x-\mu_X)^2 f_X(x) = 0$

$\Leftrightarrow (x-\mu_X)^2 f_X(x) = 0, \forall x \in \mathcal{X}$

$\Leftrightarrow (x-\mu_X)^2 = 0$, for x s.t. $f_X(x) > 0$

$\Leftrightarrow P_X(X=\mu_X) = 1$.

pmf: 1 at $c = \mu_X$ ← degenerate

cf. Thm in LNp.5-16

➢ Theorem. For $a, b \in \mathbb{R}$, $Var(aX+b) = a^2 Var(X)$ (gone) ⇒ $\sigma_{ax+b} = |a| \sigma_X$

(a, b: fixed constants)

proof. Let $Y = aX + b$, then $E(Y) = a \cdot \mu_X + b \equiv \mu_Y$.

$$Var(Y) = E(Y-\mu_Y)^2 = E[(aX+b) - (a\mu_X + b)]^2$$
$$= E[a^2(X-\mu_X)^2] = a^2 E(X-\mu_X)^2 = a^2 Var(X)$$

$a>1$, $0<a<1$, $a<0$



➢ Theorem. If X is a (discrete) r.v. with mean μ_X, then for any $c \in \mathbb{R}$,

$$E[(X-c)^2] = \sigma_X^2 + (c-\mu_X)^2 \quad \text{-(*)}$$

mean square error = Var + bias²

$(X-c)$: deviation of X from c



proof.

1. X: given, c: vary ⇒ predict X
2. c: fixed (unknown), X: vary ⇒ estimate c

$$E[(X-c)^2] = E[(X-\mu_X+\mu_X-c)^2] = \sum_{x\in\mathcal{X}}[(x-\mu_X+\mu_X-c)^2]f_X(x)$$

(X: random; μ_X, c: fixed)

$$= \sum_{x\in\mathcal{X}}[(x-\mu_X)^2 + 2(x-\mu_X)(\mu_X-c) + (\mu_X-c)^2]f_X(x)$$

$$= \sum_{x\in\mathcal{X}}(x-\mu_X)^2 f_X(x) + 2(\mu_X-c)\sum_{x\in\mathcal{X}}(x-\mu_X)f_X(x) + (\mu_X-c)^2\sum_{x\in\mathcal{X}}f_X(x)$$

$\sum_{x\in\mathcal{X}}(x-\mu_X)^2 f_X(x) = \sigma_X^2$; $\sum_{x\in\mathcal{X}}(x-\mu_X)f_X(x) = 0$ (LNp.5-14); $\sum_{x\in\mathcal{X}} f_X(x) = 1$

a 2nd-order polynomial of c → $g(c)$

- Corollary. $E[(X-c)^2]$ is minimized by letting $c=\mu_X$; and the minimum value is σ_X^2. ← (proof. It's clear by (*))

1. alternative definition of mean
2. useful for prediction

- Corollary. $\sigma_X^2 = E(X^2) - (E(X))^2$. ← (proof. Let the c in (*) be zero)

useful for calculating σ_X^2

(Recall: $E(X^2) = \sum_{x\in\mathcal{X}} x^2 f_X(x)$.)

  - Example (Committees, LNp.5-17). $Var(X) = 14/3 - 2^2 = 2/3$.

➢ $E(X^n)$ is often called the n^{th} *moment* of X ($E(X^n)$: a fixed constant)

動差 → Recall. mgf in LNp.5-5

Variance = (2nd moment) − (1st moment)²; $E(X^2)$; $E(X)$

❖ **Reading**: textbook, Sec 4.3, 4.4, 4.5

Some Commonly Used Discrete Distributions

- Bernoulli and binomial Distributions

伯努利 二項

➢ Experiment: A basic experiment with sample space Ω_0 (and p.m. P_0) is repeated n times. (random outcomes)

$\mathcal{F}_0 = \{E : E \subset \Omega_0\}$ ← cf.



In general, P_0 is not enough to uniquely determine P.

  - Example. (a) Sampling with replacement
    (b) Coin Tossing
    (c) Roulette

  - The sample space for the n trials is

$P \to \mathcal{F} \leftarrow \Omega = \Omega_0 \times \cdots \times \Omega_0 = \Omega_0^n$ (like a vector space)

$\Omega = \{\omega = (\omega_1, \cdots, \omega_n) \mid \omega_i \in \Omega_0\}$

  - **Assume** that events depending on different trials are independent ($\in \Omega$)



e.g. ith trial, $A \subset \Omega_0 \Rightarrow A' = \Omega_0 \times \cdots \times \Omega_0 \times A \times \Omega_0 \times \cdots \times \Omega_0 \subset \Omega$ (ith trial)

jth trial, $B \subset \Omega_0 \Rightarrow B' = \Omega_0 \times \cdots \times \Omega_0 \times B \times \Omega_0 \times \cdots \times \Omega_0 \subset \Omega$ ($i \neq j$)

$P(A' \cap B') = P(A')P(B') = P_0(A)P_0(B)$ ← P_0 defines P

e.g. $1 \le i_1 < i_2 < \cdots < i_k \le n$

A': formed by i_1, i_3, ... i_5 trials

B': : : i_2, i_6 trials

mutually independent.